

GENERAL INDEX TO VOLUME XII

New scientific names of plants and the final members of new combinations are printed in **bold face** type; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published scientific names and all other matter, in ordinary type.

A

admirabilis (Peniophora), 304
Aegerita candida, 226
Aegerita (Peniophora), 226
Aegerita (Sclerotium), 226
aemulans (Peniophora), 308
affinis (Peniophora), 266
Aglaonema Treubii, plastids of, 156, 175, 198
alba (Peniophora), 297
albofarcta (Peniophora), 228
albo-straminea (Peniophora), 305
albugo (Peniophora), 231
albula (Peniophora), 231
Allescheri (Gloeopeniophora), 301
Allescheri (Kneiffia), 301
Allescheri (Peniophora), 301
alutaria (Peniophora), 332
 Amide-nitrogen, 369
 Amino-nitrogen, 369
 Ammonia-nitrogen, 369
amoena (Peniophora), 276
 Apparatus used in nitrogen determination experiments, for the evaporation of samples under vacuum, 380
 Apple, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
 Apple scab, effect of sulphur treatment on, 136, 412
arachnoides (Peniophora), 220
argentea (Peniophora), 346
Arisaema triphyllum, plastids of, 157, 175, 202
 Armstrong, Joanne Karrer, B. M. Dugar and. The effect of treating the virus of tobacco mosaic with the juices of various plants, 359
 Asparagine, 369
Asparagus officinalis, crystals obtained by addition of alcoholic potash, 206; plastids of, 156, 170, 200
Aspergillus niger, effect of pokeweed juices on, 363
asperipilata (Peniophora), 230
Asterostromella epiphylla, 241
aurantiaca (Gloeopeniophora), 311
aurantiaca (Kneiffia), 311
aurantiaca (Peniophora), 310
aurantiacum (Corticium), 311

B

Bacterium prodigiosum, effect of pokeweed juice on, 363
borealis (Peniophora), 295
Botrytis Allii, toxicity of certain sulphur suspensions to, 403
Botrytis cinerea, effect of certain forms of colloidal sulphur on germination of, 389; of precipitated lime sulphur solutions on germination of, 394; toxicity of certain sulphur suspensions to, 403
Brunella aequinoctialis, 124
Burkei (Peniophora), 282
 Burt, E. A. The Thelephoraceae of North America. XIV, 213
Burtii (Peniophora), 278

C

caesia (Peniophora), 353
caesium (Corticium), 353
cana (Peniophora), 227
canadensis (Peniophora), 260
candida (Aegerita), 226
candida (Peniophora), 226
calotrichum (Corticium), 254
Capsicum annuum, change in shape of plastids during ripening, 154, 196; color changes during ripening, 188; crystals obtained by addition of alcoholic potash, 206; pigment in, 167, 175
carnea (Peniophora), 354
carneum (Corticium), 354
carnea (Peniophora), 325
 Carotin, 178, 184
 Carotinoids in fruits, pigment studies with special reference to, 145
 Carrot pigments, 184
Celastrus scandens, 157; crystals obtained by addition of alcoholic potash, 206; plastids of, 170, 202
 Cherries, results of sulphur experiments on, 141
cinerea § *Corticium* (Thelephora), 348
cinerea (Kneiffia), 348
cinerea (Peniophora), 348

cinereum (*Corticium*), 348
citrinella (*Peniophora*), 327
citrinellum (*Corticium*), 327
Citrullus vulgaris, pigment of, 166, 175; plastids of, 157, 204
coccineo-fulva (*Peniophora*), 253
coccineo-fulva (*Phlebia*), 253
Colletotrichum Gossypii, effect of certain forms of colloidal sulphur on germination of, 389; toxicity of certain sulphur suspensions to, 403
 Colloidal sulphur: as a spray material, 133; compared with other sprays in New York, 137; effect of drying and aerating upon toxicity of, 409; influence of temperature upon toxicity of, 410; preparation and toxicity of, 381; practical application of, 411; toxicity of volatile products of, 408
colorea (*Peniophora*), 343
 Colorimetric method for the determination of relative amounts of pigment in tomatoes, 160
convolvens (*Peniophora*), 355
Corticium aurantiacum, 311; *caesium*, 353; *calceum*, 282; *calotrichum*, 254; *carneum*, 354; *cinereum*, 348; *citrinellum*, 327; *crocicreas*, 322; *Eichlerianum*, 261; *filamentosum*, 320; *fumigatum*, 348; *giganteum*, 216; *glabrum*, 274; *incarnatum*, 308; *inconspicuum*, 221; *laeve*, 257; *laevigatum*, 338; *martianum*, 330; *Mollerianum*, 270; *mutatum*, 299; *nudum*, 345; *ochroleucum*, 286; *pertense*, 315; *Petersii*, 274, 320; *puberum*, 313; *radicatum*, 320; *rhodellum*, 254; *rimosis-simum*, 341; *Roumeguerii*, 270; *Sambuci*, 233; *sanguineum*, 274; *seriale*, 239; *seriale*, 318; *serum*, 233; *simile*, 336; *sordidum*, 280; *subulaceum*, 288; *subaurantiacum*, 322; *subincarnatum*, 329; *subsulphureum*, 329; *tenuis*, 317; *tephrum*, 339; *velutinum*, 264; *violaceo-lividum*, 347; *viticolum*, 322
 Cotton, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
crassa (*Peniophora*), 286
Crataegus phaenopyrum, crystals obtained by addition of glycerine potash, 206; plastids from ripe fruit, 157, 202
cremea (*Kneiffia*), 261
cremea (*Peniophora*), 261
crocicreas (*Corticium*), 322

D

Daucus Carota, 208; color pigment in, 171; natural carotin crystals of, 206
decorticans (*Peniophora*), 344
 Determination of total nitrogen in plants and plant solutions, 367

Devarda method of nitrogen determination, 367
disciformis var. *borealis* (*Peniophora*), 295
Dracaena Godseffiana, plastids of, 157, 198
 Duggar, B. M., and Joanne Karrer Armstrong. The effect of treating the virus of tobacco mosaic with the juices of various plants, 359
duplex (*Peniophora*), 298

E

Eichlerianum (*Corticium*), 261
epiphylla (*Asterostromella*), 241
 Epling, Carl Clawson. Monograph of the genus *Monardella*, 1; Studies on South American Labiatae. I, 107
Evonymus americana, crystals of, 169; obtained by addition of alcoholic potash, 210; plastids of, 157; *europaea*, crystals of, obtained by addition of alcoholic potash, 210; plastids of, 157, 198, 204
exigua (*Peniophora*), 224
exilis (*Peniophora*), 239

F

farinosa (*Kneiffia*), 226
Farlowii (*Peniophora*), 343
filamentosa (*Peniophora*), 320
filamentosum (*Corticium*), 320
firma (*Peniophora*), 276
flammea (*Peniophora*), 252
flavido-alba (*Peniophora*), 248
fumigatum (*Corticium*), 348
 Fungicide, sulphur as a, 381
fusca (*Peniophora*), 244
fusco-marginata (*Peniophora*), 335

G

galochrona (*Peniophora*), 222
 Geranium, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
gigantea (*Peniophora*), 216
gigantea (*Thelophora*), 216
giganteum (*Corticium*), 216
gigaspora (*Peniophora*), 356
gilvula (*Peniophora*), 245
glabrum (*Corticium*), 274
glebulosa (*Peniophora*), 282
globifera (*Peniophora*), 219
Gloeocystidium, 213; *guttuliferum*, 247; *praetermissum*, 316; *tenuis*, 317
Gloeopeniophora, 213; *Allescheri*, 301; *aurantiaca*, 311; *incarnata*, 308
Glomerella cingulata, effect of certain forms of colloidal sulphur on germination of, 389; effect of lime-sulphur solutions on germination of, 394

Grapes, results of sulphur experiments on, 142

guttulifera (Peniophora), 247
guttuliferum (Gloeocystidium), 247
 Glycine, 369

H

heterocystidia (Peniophora), 293
hiulca (Peniophora), 272
 Howard, Grace E. Pigment studies with special reference to carotinoids in fruits, 145
humifaciens (Peniophora), 225
 Hydrogen-ion concentration: of various juices, effect of, on tobacco mosaic virus, 362; effect of, upon the stability of colloidal sulphur solution, 418; toxicity of certain forms of colloidal sulphur in relation to, 398, 399, 400, 401, 402; influence of, upon stability of colloidal sulphur suspensions, 406
 Hydrogen sulphide and sulphur dioxide, preparation of colloidal sulphur by means of, 395
Hypochnus longisporus, 229; *Sambuci*, 233; *sulphurinus*, 324

I

incarnata (Gloeopeniophora), 308
incarnata (Kneiffia), 308
incarnata (Peniophora), 307
incarnata (Thelephora), 308
incarnatum (Corticium), 308
inconspicua (Peniophora), 221
inconspicuum (Corticium), 221
inflata (Peniophora), 267
investiens (Peniophora), 307
 Irish potato, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
irregularis (Peniophora), 228
isabellina (Peniophora), 253

J

Jimson weed, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
Juniperi (Xerocarpus), 338

K

Karstenii (Peniophora), 254, 286
Karstenii (Stereum), 286
Kauffmanii (Peniophora), 296
Kneiffia, 213; *Allescheri*, 301; *aurantiaca*, 311; *cinerea*, 348; *cremea*, 261; *fari-nosa*, 226; *incarnata*, 308; *laevigata*, 338; *laevis*, 257; *longispora*, 229; *pubera* var. *villosa*, 313; *Roumeguerii*, 270; *sanguinea*, 274; *serialis*, 318; *sub-alutacea*, 288; *tenuis*, 317; *velutina*, 264

L

laeve (Corticium), 257
laevigata (Kneiffia), 338
laevigata (Peniophora), 338
laevigata (Thelephora), 338
laevigatum (Corticium), 338
laevis (Kneiffia), 257
laevis (Peniophora), 257
laevis (Thelephora), 257
laminata (Peniophora), 246
Lamium, 125; *amplexicaule*, 125
laxa (Peniophora), 224
Leonotis, 128; *nepetaefolia*, 128
Leonurus, 126; *sibiricus*, 127
lepida (Peniophora), 295
leprosa (Peniophora), 254
lilacina (Peniophora), 348
lilacina (Thelephora), 348
 Lime sulphur, 134; preparation of colloidal sulphur from, 390; toxicity of, 394
limonia (Peniophora), 275
livida (Peniophora), 239
longispora (Kneiffia), 229
longispora (Peniophora), 229
longisporus (Hypochnus), 229
Lonicera sp., color pigment in, 170; plastids of, 157, 204, 206
ludoviciana (Peniophora), 244
Luteins, 170
Lycium halimifolium, color changes during ripening, 187; plastids of, 156, 200, 206
Lycopersicin, 159, 175, 178

M

Macranthae, 33
Macrosporium sarcinaeforme, effect of certain forms of colloidal sulphur on germination of, 389, of precipitated lime sulphur on germination of, 394; toxicity of certain sulphur suspensions to, 404
Madronella, 32; *amabilis*, 50; *anemonoides*, 72; *Breveri*, 85; *candicans*, 87; *coriacea*, 50; *dentaia*, 50; *discolor*, 60; *Douglasii*, 78; *epilobioides*, 74; *exilis*, 89; *franciscana*, 48; *gigantea*, 48; *glauca*, 62; *globosa*, 46; *hypoleuca*, 53; *ingrata*, 62; *involutrata*, 46; *lanceolata*, 79, var. *glandulifera*, 84; *ledifolia*, 55; *leucocephala*, 90; *linoides*, 72; *macrantha*, 35; *modocensis*, 62; *mollis*, 48; *muriculata*, 69; *nana*, 36; *neglecta*, 52; *nervosa*, 60; *oblonga*, 72; *oblongifolia*, 62; *odoratissima*, 59, 60; *ovata*, 62; *pallida*, 66; *Palmeri*, 41; *parviflora*, 69; *parvifolia*, 69; *peninsularis*, 84; *pinetorum*, 68; *Pringlei*, 86; *purpurea*, 62; *rubella*, 62; *sanguinea*, 79; *sessilifolia*, 62; *sub-*

serrata, 48; *thymifolia*, 44; *undulata*, 76; *villosa*, 46; *viminea*, 72; *viridis*, 55
magnahypa (Peniophora), 238
Marrubium, 122; *hamatum*, 122; *vulgare*, 122
martiana (Peniophora), 330
martianum (Corticium), 330
medioburiensis (Peniophora), 328
mexicana (Peniophora), 243
miniata (Peniophora), 277
miniata (Thelephora), 277
Molleriana (Peniophora), 270
Mollerianum (Corticium), 270
Momordica, color pigment in, 170
Monarda fistulosa, 2; *punctata*, 7
Monardella, monograph of the genus, 1; diagram of relationships in, 14
Monardella acuta, 79; *anemonoides*, 72; *australis*, 70; *Breweri*, 84, 104; *candicans*, 87, 106; *candicans*, 78, 106, var. *exilis*, 89, var. *venosa*, 78; *Caroliniana*, 2; *cinerea*, 56, 98; *coriacea*, 50; *crispa*, 77; *dentata*, 50; *discolor*, 60; *Douglasii*, 2, 78, 104, 106; *Elmeri*, 85; *epilobioides*, 74, var. *erecta*, 74; *exilis*, 92, 106; *franciscana*, 48; *glabra*, 59; *glauca*, 62; *globosa*, 46; *hypoleuca*, 53, 100; *ingrata*, 62; *involuta*, 46; *lanata*, 54, 100; *lanceolata*, 79, 104, var. *microcephala*, 84; *lanceolata* var. *glandulifera*, 84; *ledifolia*, 55; *leucocephala*, 90, 106; *linoides*, 71, 102, subsp. *eulinoides*, 72, subsp. *stricta*, 74; *linoides*, 72, var. *stricta*, 74; *macrantha*, 34, 96, var. *arida*, 38, var. *Hallii*, 35, var. *tenuiflora*, 37, subsp. *eumacrantha*, 35, subsp. *nana*, 36; *macrantha*, 35, var. *longiloba*, 36, var. *nana*, 36, var. *pinetorum*, 37, var. *tenuiflora*, 35; *modocensis*, 62; *mollis*, 48; *montana*, 2; *muriculata*, 69; *nana*, 36, var. *arida*, 38, var. *leptosiphon*, 36, var. *tenuiflora*, 37; *neglecta*, 52; *nervosa*, 60; *oblonga*, 72; *odoratissima*, 57, 102, subsp. *australis*, 70, 94, 102, subsp. *discolor*, 60, 94, 102, subsp. *euodora-tissima*, 59, 94, 102, subsp. *glauca*, 62, 94, 102, subsp. *pallida*, 62, 94, 102, subsp. *parvifolia*, 69, 94, 102, subsp. *pinetorum*, 68, 94; *odoratissima*, 59; *ovata*, 62; *pallida*, 66; *Palmeri*, 41, 96; *parviflora*, 69; *parvifolia*, 69; *peninsularis*, 84; *pinetorum*, 68; *Pringlei*, 86, 106; *purpurea*, 62; *reflexa*, 50; *robusta*, 53; *rubella*, 62; *sanguinea*, 79; *saxicola*, 55, 100; *Sheltoni*, 50; *subserrata*, 48; *thymifolia*, 44, 98; *tomentosa*, 48; *tenuiflora*, 37; *undulata*, 76, 104, var. *crispa*, 77; *villosa*, 44, 48, 98, subsp. *euvillosa*, 46, var. *franciscana*, 48, subsp. *neglecta*, 52, subsp. *Sheltoni*, 50, subsp. *subserrata*, 48; *villosa* var.

glabella, 50, var. *leptosiphon*, 36, 48; *viminea*, 72; *viridis*, 55, 100
montana (Peniophora), 237
Mosaic, tobacco, the effect of treating the virus of, with the juices of various plants, 359
mutata (Peniophora), 299
mutatum (Corticium), 299

N

Nitrate-nitrogen, 369
 Nitrogen, determination of total, in plants and plant solutions, 367
nuda (Peniophora), 345
nuda (Thelephora), 345
nudum (Corticium), 345

O

ochracea (Peniophora), 345
ochroleucum (Corticium), 286
odontioides (Peniophora), 223
odorata (Peniophora), 289
odorata (Phanerochaete), 286, 289
odoratum (Stereum), 289

P

Peaches, results of sulphur experiments on, 141
Peckii (Peniophora), 291
 Peniophora, 213; *admirabilis*, 304; *Aegeria*, 226; *aemulans*, 308; *affinis*, 266; *alba*, 297; *albofarcta*, 228; *albostraminea*, 305; *albugo*, 231; *albula*, 231; *Allescheri*, 301; *alutaria*, 332; *amoena*, 276; *arachnoidea*, 220; *argentea*, 346; *asperipilata*, 230; *aurantiaca*, 310; *borealis*, 295; *Burkei*, 282; *Burtii*, 278; *caesia*, 353; *cana*, 227; *canadensis*, 260; *candida*, 226; *carnea*, 354; *carnosa*, 325; *cinerea*, 348; *citrinella*, 327; *coccineofulva*, 253; *colorea*, 343; *convolvens*, 355; *crassa*, 286; *cremea*, 261; *decorticans*, 344; *disciformis* var. *borealis*, 295; *duplex*, 298; *exigua*, 224; *exilis*, 239; *Farlowii*, 343; *filamentosa*, 320; *firma*, 276; *flammea*, 252; *flavido-alba*, 248; *fusca*, 244; *fusco-marginata*, 335; *galochroa*, 222; *gigantea*, 216; *gigaspora*, 356; *gilvidula*, 245; *glebulosa*, 282; *globifera*, 219; *guttulifera*, 247; *heterocystidia*, 293; *hiulca*, 272; *humifaciens*, 225; *incarnata*, 307; *inconspicua*, 221; *inflata*, 267; *investiens*, 307; *irregularis*, 228; *isabellina*, 253; *Karstenii*, 254, 286; *Kauffmanii*, 296; *laevigata*, 338; *laevis*, 257; *laminata*, 246; *laxa*, 224; *lepida*, 295; *leprosa*, 254; *lilacina*, 348;

limonia, 275; **livida**, 239; **longispora**, 229; **ludoviciana**, 244; **magna-hypha**, 238; **martiana**, 330; **medio-buriensis**, 328; **mexicana**, 243; **miniata**, 277; **Molleriana**, 270; **montana**, 237; **mutata**, 299; **nuda**, 345; **ochracea**, 345; **odontioides**, 223; **odorata**, 289; **Peckii**, 291; **pertenuis**, 315; **phosphorescens**, 273; **phyllophila**, 241; **piliseta**, 242; **pilosa**, 291; **praetermissa**, 316; **pruinata**, 340; **pubera**, 313; **pubera f. villosa**, 313; **Ravenelii**, 269; **rhodella**, 254; **rhodochroa**, 254; **rimosissima**, 341; **Roumeguerii**, 270; **Sacchari**, 328; **Sambuci**, 233; **sanguinea**, 274; **separans**, 332; **septocystidia**, 260; **serialis**, 318; **serialis**, 239; **Seymouriana**, 337; **Sheari**, 268; **similis**, 336; **sordida**, 280; **stratosa**, 333; **subalutacea**, 288; **subapiculata**, 280; **subcremea**, 303; **subiculosa**, 259; **subsulphurea**, 329; **sulphurina**, 324; **tabacina**, 334; **Taxodii**, 306; **tenella**, 298; **tenuis**, 317; **tephra**, 339; **terricola**, 237; **texana**, 251; **Thujae**, 236; **typhicola**, 319; **unicolor**, 320; **velutina**, 264; **verniciosa**, 250; **versata**, 305; **verticillata**, 285; **violaceo-livida**, 347; **viticola**, 322; **Weirii**, 342; **zonata**, 245
pertenuis (*Corticium*), 315
pertenuis (*Peniophora*), 315
Petersii (*Corticium*), 274, 320
Phanerochaete odorata, 286, 289
phosphorescens (*Peniophora*), 273
Phlebia coccineo-fulva, 253
phyllophila (*Peniophora*), 241
Pigment studies with special reference to carotinoids in fruits, 145
piliseta (*Peniophora*), 242
pilosa (*Peniophora*), 291
Plums, results of sulphur experiments on, 141
Pokeweed, effect of juices of, on pathogenicity of tobacco mosaic virus, 359
praetermissa (*Peniophora*), 316
praetermissum (*Gloeocystidium*), 316
pruinata (*Peniophora*), 340
pruinatum (*Stereum*), 340
Prunella, 124; *vulgaris*, 124
pubera (*Peniophora*), 313
pubera (*Thelephora*), 313
pubera var. villosa (*Kneiffia*), 313
pubera f. villosa (*Peniophora*), 313
puberum (*Corticium*), 313
Pycnanthae, 33, 42

R

radicatum (*Corticium*), 320
Ranker, Emery R. Determination of total nitrogen in plants and plant

solutions: a comparison of methods with modifications, 367
Ravenelii (*Peniophora*), 269
rhodella (*Peniophora*), 254
rhodellum (*Corticium*), 254
rhodochroa (*Peniophora*), 254
Rhus canadensis, color pigment in, 175; crystals obtained by addition of alcoholic potash, 206
rimosissima (*Peniophora*), 341
rimosissimum (*Corticium*), 341
Rosa rugosa, color pigment in, 168; plastids of, 155, 196
Rosmarinus, 120; *chilensis*, 120; *officinalis*, 120
Roumeguerii (*Corticium*), 270
Roumeguerii (*Kneiffia*), 270
Roumeguerii (*Peniophora*), 270

S

Sacchari (*Peniophora*), 328
Salicylic-thiosulphate method of nitrogen determination, 367; modification of, 368
Sambuci (*Corticium*), 233
Sambuci (*Hypochnus*), 233
Sambuci (*Peniophora*), 233
sanguinea (*Kneiffia*), 274
sanguinea (*Peniophora*), 274
sanguinea (*Thelephora*), 274
sanguineum (*Corticium*), 274
Sclerotinia cinerea, effect of certain forms of colloidal sulphur on germination of, 389; of lime-sulphur solutions on germination of, 394; toxicity of certain sulphur suspensions to, 404
Sclerotium Aegeria, 226
separans (*Peniophora*), 332
septocystidia (*Peniophora*), 260
sera (*Thelephora*), 233
seriale (*Corticium*), 318
serialis (*Kneiffia*), 318
serialis (*Peniophora*), 318
serum (*Corticium*), 233
Seymouriana (*Peniophora*), 337
Sheari (*Peniophora*), 268
simile (*Corticium*), 336
similis (*Peniophora*), 336
Sodium thiosulphate and sulphuric acid, preparation of colloidal sulphur from, 383
Solanum Dulcamara, plastids of, 156, 198; crystals of, 169, obtained by addition of alcoholic potash, 206, 210
Solanum Lycopersicum, crystals in ripe fruit of, 210; plastids of, 158, 198; pigment in, 168
Solanum Pseudo-capsicum, plastids of, 156, 202
Sorbus sitchensis, crystals obtained by addition of alcoholic potash, 206; plastids of, 156, 202; pigment in, 169

sordida (Peniophora), 280
sordidum (Corticium), 280
 Spectroscopic analysis of fruits, 176
 Spray material, colloidal sulphur as a, 133, 382
 Squash, effect of juices of, on pathogenicity of tobacco mosaic virus, 362
Stereum Karstenii, 286; *odoratum*, 289; *pruinatum*, 340; *umbrinum*, 341
stratosa (Peniophora), 333
subalutacea (Kneiffia), 288
subalutacea (Peniophora), 288
subalutaceum (Corticium), 288
subapiculata (Peniophora), 280
subawantiacum (Corticium), 322
subermea (Peniophora), 303
subiculososa (Peniophora), 259
subincarnatum (Corticium), 329
subsulphurea (Peniophora), 329
subsulphureum (Corticium), 329
subsulphureus (Xerocarpus), 338
 Sulphur: as a fungicide, 381; colloidal, as a spray material, 133, 382; precipitated, preparation of, 134
 Sulphur dioxide and hydrogen sulphide, preparation of colloidal sulphur by means of, 395
 Sulphuric acid and sodium thiosulphate, preparation of colloidal sulphur from, 383
sulphurina (Peniophora), 324
sulphurina (Tomentella), 324
sulphurinus (Hypochnus), 324
 Sweet potato, effect of juices of, on pathogenicity of tobacco mosaic virus, 362

T

tabacina (Peniophora), 334
Taxodii (Peniophora), 306
 Temperature, influence of, upon the toxicity of colloidal sulphur, 410; relation of, to ripening of fruits, 186
tenella (Peniophora), 298
tenua (Corticium), 317
tenua (Gloeocystidium), 317
tenuis (Kneiffia), 317
tenuis (Peniophora), 317
tephra (Peniophora), 339
tephrum (Corticium), 339
terricola (Peniophora), 237
Teucrium, 111; *bicolor*, 115; *Cavenellaei*, 115; *Chamaedrifolium*, 112; *chilense*, 115; *cubense*, 111, subsp. *Chamaedrifolium*, 112, subsp. *cordobense*, 113, 130, var. *leucanthum*, 114; *cubense*, 113; *Grisebachii*, 113; *heterophyllum*, 115; *hirtum*, 118; *inflatum*, 116; *laevigatum*, 112; *leucanthum*, 114; *nudicaule*, 113; *orchideum*, 115; *palustre*, 118; *tenuipes*, 119, 133; *tripartitum*, 113; *vesicarium*, 116

texana (Peniophora), 251
Thelephora cinerea § *Corticium*, 348; *gigantea*, 216; *incarnata*, 308; *laevigata*, 338; *laevis*, 257; *lilacina*, 348; *miniata*, 277; *nuda*, 345; *pubera*, 313; *Sambuci*, 233; *sanguinea*, 274; *sera*, 233; *velutina*, 264; *violaceo-livida*, 347; *viticola*, 322
 Thelephoraceae of North America, The, XIV, 213
Thujae (Peniophora), 236
 Tisdale, L. E. Colloidal sulphur: preparation and toxicity, 381
 Tobacco mosaic, the effect of treating the virus of, with the juices of various plants, 359
 Tomatoes, colorimetric method for determination of relative amounts of pigment in, 160
Tomentella sulphurina, 324
typhicola (Peniophora), 319

U

umbrinum (Stereum), 341
unicolor (Peniophora), 320
 Ustilago Hordei, effect of lime sulphur solutions on germination of, 394; toxicity of certain sulphur suspensions to, 404

V

velutina (Kneiffia), 264
velutina (Peniophora), 264
velutina (Thelephora), 264
velutinum (Corticium), 264
vernica (Peniophora), 250
versata (Peniophora), 305
verticillata (Peniophora), 285
Viburnum Opulus, plastids of, 157, 202; pigment in, 170
violaceo-livida (Peniophora), 347
violaceo-livida (Thelephora), 347
violaceo-lividum (Corticium), 347
viticola (Peniophora), 322
viticola (Thelephora), 322
viticolum (Corticium), 322

W

Weiri (Peniophora), 342

X

Xanthophyll, 173, 178
Xerocarpus Juniperi, 338; *subsulphureus*, 329

Y

Young, H. C. Colloidal sulphur as a spray material, 133

Z

zonata (Peniophora), 245

